

## Synthesis, Characterization, and Antibacterial Evaluation of New Vanillic Acid Derivatives

Mostafa F. Tawfeeq<sup>\*1</sup> and Ahlam J. Qassir<sup>\*\*</sup>

<sup>\*</sup> Department of Pharmacy, College of Pharmacy, University of Tikrit, Salahuddin, Iraq

<sup>\*\*</sup> Department of Pharmaceutical Chemistry, College of Pharmacy, University of Baghdad, Baghdad, Iraq

### Abstract

Hydrazide Schiff bases (hydrazones) and 2,5-disubstituted-1,3,4-oxadiazole derivatives exhibit diverse biological activities that include antibacterial, antifungal, antitubercular, antiviral, anticancer, anti-inflammatory, and analgesia; so that new derivatives, **compounds (5-8)** of vanillic acid based on 1,3,4-oxadiazole as scaffold unit were synthesized through multi-steps, and characterized by thin layer chromatography and spectroscopically by Fourier-transform Infrared (FTIR) and Proton nuclear magnetic resonance (<sup>1</sup>HNMR). **Compounds (5-8)** were evaluated for their antibacterial activity by the disk diffusion method. **Compounds (5-8)** were showed moderate and comparable to the activities of amoxicillin and isoniazid against *Escherichia coli* (*E. coli*), but less than that of cefixime and nitrofurantoin which their activities were high. **Compound (6)** and **(7)** had shown moderate and comparable to the activities of amoxicillin and cefixime against *Klebsiella pneumoniae* (*K. pneumoniae*), but less than that of nitrofurantoin which its activity was high. **Compound (6)** and **(7)** had shown moderate and comparable to the activity of amoxicillin against *Staphylococcus aureus* (*S. aureus*), while the activities of cefixime and nitrofurantoin were high. **Compound (6)** was moderately active against *Bacillus subtilis* (*B. subtilis*), while the activities of amoxicillin, cefixime, and nitrofurantoin were high.

**Keywords:** Hydrazide, Schiff Base, Oxadiazole, Antibacterial.

تصنيع ، تشخيص ، وتقييم الفعالية المضادة للبكتريا لمشتقات جديدة لحمض الفانيلين  
مصطفى فايز توفيق<sup>\*1</sup> و احلام جميل قصير<sup>\*\*</sup>

فرع الكيمياء الصيدلانية، كلية الصيدلة، جامعة تكريت، صلاح الدين، العراق .

فرع الكيمياء الصيدلانية، كلية الصيدلة، جامعة بغداد، بغداد، العراق .

### الخلاصة

قواعد شف و الحلقة الأروماتية الغير متجانسة ٤، ٣، ١-اوكساديزول المعوضة في الموقعين ٢ و ٥ لهما العديد من الفعاليات الحيوية كأن تكون مضادات للبكتيريا والفطريات وبكتريا التدرن والفايروسات والخلايا السرطانية أو أن تكون مضادات للالتهابات أو تعمل كمسكنات للألم. لذلك تم تصنيع مشتقات جديدة لحمض الفانيلين المركبات (٥-٨) بالاعتماد على الحلقة الأروماتية الغير متجانسة ٤، ٣، ١-اوكساديزول كوحدة بناء رئيسية بعدة خطوات وهذه المركبات تم تشخيصها عن طريق الكروماتوغرافيا (استشراب الطبقة الرقيقة) و عن طريق مطياف الأشعة تحت الحمراء، الرنين النووي المغناطيسي للبروتون. **المركبات (٥-٨)** تم تقييم فعاليتها المضادة للبكتريا باستخدام طريقة الانتشار. **المركبات (٥-٨)** أظهرت فعالية متوسطة يمكن مقارنتها بفعالية الاموكسلين والايرونيازيد ضد **البكتريا الاشريكية القولونية** ولكنها كانت أقل من فعالية السفيكزيم والنيتروفورانتوين حيث كانت فعاليتها عالية. **المركبان (٦)** و **(٧)** أظهرت فعالية متوسطة يمكن مقارنتها بفعالية الاموكسلين والسفيكزيم ضد **البكتريا الكلبسيلا الرئوية** ولكنها كانت أقل من فعالية النيتروفورانتوين حيث كانت له فعالية عالية. **المركبان (٦)** و **(٧)** أظهرت فعالية متوسطة يمكن مقارنتها بفعالية الاموكسلين ضد **البكتيريا الكروية العنقودية الذهبية** بينما السفيكزيم والنيتروفورانتوين كانت لهما فعالية عالية. **المركب (٦)** كانت له فعالية متوسطة ضد **البكتريا العنقودية الرقيقة** بينما فعالية الاموكسلين والسفيكزيم والنيتروفورانتوين كانت فعالية عالية. الكلمات المفتاحية: هيدرازيد، قواعد شف، اوكساديزول، مضادات بكتيرية.

### Introduction

Heterocyclic compounds are used in many biological fields, due to their different activities, and are considered as one of the principal classes of organic compounds, that are used in the development of several pharmaceutically essential compounds<sup>(1,2)</sup>.

Oxadiazoles are important five-membered aromatic heterocyclic containing oxygen and two nitrogens in their structure. Because the oxadiazole ring is structurally rigid, various functional groups are easily introduced into the ring. Valuable biological activities are associated with oxadiazole derivatives<sup>(3)</sup>, such as antitumor<sup>(4)</sup>, anti-inflammatory<sup>(5)</sup>,

antimicrobial<sup>(6)</sup>, antifungal<sup>(7)</sup>, and anticonvulsant<sup>(8)</sup>.

In synthetic medicinal chemistry, to improve the biological activity of new drugs with respect to the corresponding lead compounds, hybridization-combination of different pharmacophores in one structure-is one of the techniques being followed.<sup>(9,10)</sup> Hydrazide Schiff base derivatives (hydrazones) are good scaffolds for various pharmaceutical applications, and characterized by the presence of highly reactive azomethine group (-CO-NH-N=CH-).<sup>(11)</sup> Their biological activities include antibacterial<sup>(12)</sup>, antifungal<sup>(13)</sup>, antitubercular<sup>(14)</sup>, antiviral<sup>(15)</sup>, anticancer<sup>(16)</sup>, anti-inflammatory<sup>(17)</sup>, and analgesia.<sup>(18)</sup>

<sup>1</sup>corresponding author E-mail: m8mostafa@gmail.com

Received:27 / 1 / 2020

Accepted:20 / 5 / 2020

## Material and Methods

Chemicals supplied by hyper-chem (China) were used. Aluminum sheets pre-coated with Silica gel GF<sub>254</sub> (type 60), exposed to UV-254 nm, were used for thin-layer chromatography (TLC) to monitor the completion of reactions and to test the purity of compounds. Two solvent systems (S<sub>1</sub> and S<sub>2</sub>) were used: S<sub>1</sub>(toluene:ethylacetate:ethanol (3:2:1)) and S<sub>2</sub>(ethylacetate:methanol:ammonia (5:3.5:1.5)). Melting points were incorrected and detected by using Stuart SMP3 melting point

apparatus in open capillary tubes. All synthesized derivatives were characterized by spectroscopic analysis (Fourier-transform Infrared (FTIR) which was performed at Baghdad university-college of pharmacy) and (Proton nuclear magnetic resonance (<sup>1</sup>HNMR) which was performed by Chemistry Analysis Center (CAC)) at Iraq/ Baghdad<sup>(19-22)</sup>.

### Chemical synthesis

The target compounds were synthesized by multistep(s) reactions as shown in the scheme in Figure (1).




Figure 1. Multistep(s) scheme for synthesis of targeted compounds.

### Synthesis of Methyl 4-hydroxy-3-methoxybenzoate; compound (1)<sup>(23)</sup>:

Vanillic acid (7.0g, 42mmole) was dissolved in 75ml of absolute methanol (99.8%), the temperature of this solution was lowered to 0°C by ethanol-water ice bath, 5ml of concentrated H<sub>2</sub>SO<sub>4</sub> was added dropwise, and the mixture was stirred at room

temperature for 48 hours, then refluxed for 7 hours. The reaction mixture was poured in beaker containing crashed ice (100ml), the formed precipitate was filtered, collected, and washed with 5% NaHCO<sub>3</sub> aqueous solution, filtered again, and dried by a warmed current of air, giving 7.0g of compound (1).

**Synthesis of 4-Hydroxy-3-methoxybenzohydrazide; compound (2)<sup>(24)</sup>:**

**Compound (1)**, (5.0g, 27mmole) was dissolved in a minimum amount (10ml) of absolute ethanol (99.9%), (13.5g, 270mmole) of 80% hydrazine hydrate was added gradually. The mixture was refluxed for 3-4 hours (monitored by TLC). Then the reaction mixture was cooled and precipitate begin to appear, which was filtered and dried in oven (adjusted at 60°C), giving 3.5g of **compound (2)**.

**Synthesis of 4-(5-Mercapto-1,3,4-oxadiazol-2-yl)-2-methoxyphenol; compound (3)<sup>(25)</sup>:**

To suspension of (1.25g, 6.86mmole) of **compound (2)** in 30ml of 50% aqueous ethanolic solution, (0.785g, 10.3mmole) of CS<sub>2</sub> was added with stirring for few minutes, followed by the addition of (0.7g, 10.3mmole) potassium hydroxide (KOH), and the mixture was refluxed for 12 hours (monitored by TLC, and notation the evolution of H<sub>2</sub>S gas by strip soaked with lead acetate aqueous solution; which was turned black as indication of evolution of H<sub>2</sub>S gas). The mixture poured in a beaker containing crashed ice (30 ml) and then concentrated HCl was added dropwise until pH became (2-3). The formed precipitate was filtered, dried, and purified by base-acid precipitation method (i.e.; the product was dissolved in water by the aid of equimilimole of sodium hydroxide or triethylamine, filtered, and the clear filtrate was treated with concentrated HCl which was added dropwise until the precipitate formed again), giving 0.9g of **compound (3)**.

**Synthesis of Ethyl 2-((5-(4-hydroxy-3-methoxyphenyl)-1,3,4-oxadiazol-2-yl)thio)acetate; compound (4)<sup>(26)</sup>:**

**Compound (3)**, (0.5g, 2.23mmole,) was dissolved in 2ml of absolute ethanol (99.9%), (0.155g, 1.115mmole) of anhydrous K<sub>2</sub>CO<sub>3</sub> was added with stirring, after few minutes a white precipitate was formed, (0.37241g, 2.23mmole) of ethyl 2-bromoacetate was added dropwise. The reaction mixture refluxed for 2 hours (monitored by TLC) and left for stirring gently overnight. Then distilled water (D.W) was added gradually until precipitate was formed; then filtered, dried and crystallized from minimum amount (the amount that was just covered the powder) of boiled absolute ethanol giving 0.5g of pure product; **compound (4)**.

**Synthesis of 2-((5-(4-Hydroxy-3-methoxyphenyl)-1,3,4-oxadiazol-2-yl) thio) acetohydrazide; compound (5)<sup>(27)</sup>:**

**Compound (4)**, (0.5g, 1.6167mmole) was dissolved in a 8ml of hot absolute ethanol (99.9%), (0.16167g, 3.233mmole) of 80% hydrazine hydrate was then added, after 2 hours of vigorous mixing at room temperature, precipitate was formed, and left to stir gently overnight. Then, washed with (10ml)

absolute ethanol, filtered, dried, and used without further purification.

**Synthesis of N'-(4-hydroxy-3-methoxybenzylidene)-2-((5-(4-hydroxy-3-methoxyphenyl)-1,3,4-oxadiazol-2-yl)thio)acetohydrazide; compound (6), N'-((1H-pyrrol-2-yl)methylene)-2-((5-(4-hydroxy-3-methoxyphenyl)-1,3,4-oxadiazol-2-yl)thio)acetohydrazide; compound (7), and 2-((5-(4-hydroxy-3-methoxyphenyl)-1,3,4-oxadiazol-2-yl)thio)-N'-(4-methoxybenzylidene)acetohydrazide; compound (8)<sup>(28)</sup>:**

An appropriate aldehydes; 4-hydroxy-3-methoxybenzaldehyde (**Vanillin**) (0.039g, 0.257mmole) for **compound (6)**, 1H-pyrrole-2-carbaldehyde (0.0257g, 0.257mmole) for **compound (7)**, and 4-methoxybenzaldehyde (**p-Anisaldehyde**) (0.035g, 0.257mmole) for **compound (8)**, were dissolved in 5ml of absolute methanol (99.8%) and stirred for 15-30 minutes in the presence of 2-3 drops of glacial acetic acid; then (0.075g, 0.25mmole) of **compound (5)** was added and the suspension that formed had been refluxed for 1-2 hours. The reaction mixture was left to stir overnight, then the formed precipitate was filtered, dried and crystallized from boiled methanol.

**Antibacterial assay**

Well diffusion assay was carried out through using bacterial suspension of nearly (1.5×10<sup>6</sup>CFU/ml) obtained from McFarland turbidity standard (number 0.5). This was used to inoculate by swabbing the surface of Mueller Hinton Agar (MHA) plates. The excess liquid was dried by air under a sterile hood. In each agar plate of examined bacteria, four wells were made, and (80μl) of each concentration of the synthesized compound was poured to it. The plates were incubated at 37°C for 24 hours. The evaluation of antibacterial activity was based on the measurement of the diameter of the inhibition zone formed around the well.<sup>(29)</sup>

**Results****Chemistry**

**Compound (1)** (C<sub>9</sub>H<sub>10</sub>O<sub>4</sub>): off-white powder, yield 91%, m.p: 60-62°C (reported m.p: 64°C)<sup>(30)</sup>. **The characteristic bands in FTIR spectrum in cm<sup>-1</sup>:** (1686) C=O stretching vibration band, (1277) C-O stretching vibration band of aromatic ester.

<sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) in ppm: 3.80, 3H, s, set (a) protons; 3.83, 3H, s, set (b) protons; 6.87-6.89, 1H, d, set (c) proton; 7.44-7.49, 2H, m, set (d) protons; 9.97, 1H, s, set (e) proton.

**Compound (2)** (C<sub>8</sub>H<sub>10</sub>N<sub>2</sub>O<sub>3</sub>): white powder, yield 70%, m.p: 208-210°C (reported m.p: 210-211°C)<sup>(31)</sup>.

**The characteristic bands in FTIR spectrum in cm<sup>-1</sup>:** (3310) phenolic OH stretching vibration band overlapped with NH<sub>2</sub> asymmetric stretching vibration band, (3256) NH amide stretching

vibration band, (3209) NH<sub>2</sub> symmetric stretching vibration band, (1628) stretching vibration band of carbonyl amide, (1601) NH bending vibration band of hydrazide amine, (1585) NH bending vibration band of hydrazide amide.

<sup>1</sup>HNMR(400MHz, DMSO-*d*<sub>6</sub>) in ppm: 3.82, 3H, s, set (a) protons; 4.42, 2H, s, set (b) protons; 6.81-6.83, 1H, d, set (c) proton; 7.33-7.44, 2H, m, set (d) protons; 9.56, 2H, s, set (e) protons.

**Compound (3) (C<sub>9</sub>H<sub>8</sub>N<sub>2</sub>O<sub>3</sub>S):** faint yellow powder, yield 59%, M.P: 186-189°C. **The characteristic bands in FTIR spectrum in cm<sup>-1</sup>:** (3452) phenolic OH stretching vibration band, (3155) NH broad (br) stretching vibration band, 2669 weak (w) SH stretching vibration band.

<sup>1</sup>HNMR(400MHz, DMSO-*d*<sub>6</sub>) in ppm: 3.84, 3H, s, set (a) protons; 6.92-6.94, 1H, d, set (b) protons; 7.30-7.37, 2H, m, set (c) protons; 10.04, 1H, s, set (d) proton; 14.58, 1H, s, set (e) proton.

**Compound (4) (C<sub>13</sub>H<sub>14</sub>N<sub>2</sub>O<sub>5</sub>S):** white crystalline powder, yield 65.5%, m.p: 118-120°C. **The characteristic bands in FTIR spectrum in cm<sup>-1</sup>:** (3400-2800) broad phenolic OH stretching vibration band, (1744) carbonyl stretching vibration band of aliphatic saturated ester, (1173) stretching vibration band of C-O of aliphatic saturated ester.

<sup>1</sup>HNMR(500MHz, DMSO-*d*<sub>6</sub>) in ppm: 1.18-1.21, 3H, t, set (a) protons; 3.86, 3H, s, set (b) protons; 4.14-4.18, 2H, q, set (c) protons; 4.26, 2H, s, set (d) protons; 6.95-6.96, 1H, d, set (e) proton; 7.41-7.43, 2H, m, set (f) protons; 9.97, 1H, s, set (g) proton.

**Compound (5) (C<sub>11</sub>H<sub>12</sub>N<sub>4</sub>O<sub>4</sub>S):** white powder, yield 65%, m.p: 196-199°C. **The characteristic bands in FTIR spectrum in cm<sup>-1</sup>:** (3506) stretching vibration band of phenolic OH, (3341) asymmetric NH<sub>2</sub> stretching vibration band, (3256) NH amide stretching vibration band, (3217) symmetric NH<sub>2</sub> stretching vibration band, (1682) C=O carbonyl amide stretching vibration band, (1655) NH<sub>2</sub> bending vibration band.

<sup>1</sup>HNMR(500MHz, DMSO-*d*<sub>6</sub>) in ppm: 3.86, 3H, s, set (a) protons; 4.00, 2H, s, set (b) protons; 4.36, 2H, s, set (c) protons; 6.93-6.95, 1H, d, set (d) proton; 7.41-7.44, 2H, m, set (e) protons; 9.41, 1H, s, set (f) proton; 9.93, 1H, s, set (g) proton.

**Compound (6) (C<sub>19</sub>H<sub>18</sub>N<sub>4</sub>O<sub>6</sub>S):** white powder, yield 40%, m.p: 196-200°C. **The characteristic bands in FTIR spectrum in cm<sup>-1</sup>:** (3449) two Phenolic OH<sub>s</sub> stretching vibration bands, (3182) NH secondary amide stretching vibration band, (1678) C=O amide carbonyl stretching vibration band, (1601) C=N imine stretching vibration, C=N stretching vibration of oxadiazole and C=C aromatic skeletal stretching vibration (overlapped) bands.

<sup>1</sup>HNMR(500MHz, DMSO-*d*<sub>6</sub>) in ppm: 3.81, 3.80, 3H, 2s, set (a) protons [syn/anti-syn]; 3.83, 3H, s, set (b) protons; 4.16, 4.61, 2H, 2s, set (c) protons [syn/anti-syn]; 6.80-6.83, 1H, m, set (d) proton; 6.90-6.93, 1H, m, set (e) proton; 7.07-7.09, 1H, m,

set (f) proton; 7.26-7.27, 1H, m, set (g) proton; 7.39-7.42, 2H, m, set (h) protons; 7.91, 8.08, 1H, 2s, set (i) proton [syn/anti-syn]; 9.52, 1H, s, set (j) proton; 9.94, 1H, s, set (k) proton; 11.60, 11.63, 1H, 2s, set (l) proton [cis/trans].

**Compound (7) (C<sub>16</sub>H<sub>15</sub>N<sub>5</sub>O<sub>4</sub>S):** white powder, yield 53.4%, m.p: 226-228°C. **The characteristic bands in FTIR spectrum in cm<sup>-1</sup>:** (3302) phenolic OH and NH pyrrole stretching vibration (overlapped) bands, (3209) NH amide stretching vibration band, (1663) amide carbonyl stretching vibration band, (1620) NH pyrrole bending vibration and imine stretching vibration (overlapped) bands.

<sup>1</sup>HNMR(500MHz, DMSO-*d*<sub>6</sub>) in ppm: 3.83, 3H, s, set (a) protons; 4.17, 4.60, 2H, 2s, set (b) protons [syn/anti-syn]; 6.11-6.14, 1H, m, set (c) proton; 6.44-6.49, 1H, m, set (d) proton; 6.90-6.95, 2H, m, set (e) protons; 7.40-7.43, 2H, m, set (f) protons; 7.86, 8.04, 1H, 2s, set (g) proton [syn/anti-syn]; 9.94, 1H, s, set (h) proton; 11.40, 1H, s, set (i) proton; 11.45, 11.48, 1H, 2s, set (j) proton [cis/trans].

**Compound (8) (C<sub>19</sub>H<sub>18</sub>N<sub>4</sub>O<sub>5</sub>S):** White powder, yield 72%, m.p: 190-193°C. **The characteristic bands in FTIR spectrum in cm<sup>-1</sup>:** (3495) Phenolic OH stretching vibration band, (3182) NH secondary amide stretching vibration band, (1666) C=O amide carbonyl stretching vibration band, (1609) C=N imine stretching vibration, C=N stretching vibration of oxadiazole and C=C aromatic skeletal vibration (overlapped) bands.

<sup>1</sup>HNMR(500MHz, DMSO-*d*<sub>6</sub>) in ppm: 3.79, 3.80, 3H, 2s, set (a) protons [syn/anti-syn]; 3.83, 3H, s, set (b) protons; 4.18, 4.59, 2H, 2s, set (c) protons [syn/anti-syn]; 6.90-7.01, 3H, m, set (d) protons; 7.39-7.42, 2H, m, set (e) protons; 7.61-7.65, 2H, m, set (f) protons; 7.97, 8.14, 1H, 2s, set (g) proton [syn/anti-syn]; 9.94, 1H, s, set (h) proton; 11.64, 11.69, 1H, 2s, set (i) proton [cis/trans].

#### Anti-bacterial evaluation

The antibacterial activities of the synthesized compounds; **compounds (5-8)** were evaluated against six bacteria and compared with four standard antibiotics; amoxicillin, cefixime, nitrofurantoin, and isoniazid. Dimethyl sulfoxide (DMSO) was used as solvent and control. It's evident from the data displayed in the table (1), **compound (5)** showed moderate antibacterial activity against *Escherichia coli* (*E. coli*). **Compound (6)** showed moderate activity towards *Staphylococcus aureus* (*S. aureus*), *Bacillus subtilis* (*B. subtilis*), *Escherichia coli* (*E. coli*), and *Klebsiella pneumoniae* (*K. pneumoniae*). **Compound (7)** showed moderate activity towards *Staphylococcus aureus* (*S. aureus*), *Escherichia coli* (*E. coli*), and *Klebsiella pneumoniae* (*K. pneumoniae*). **Compound (8)** was slightly active against *Staphylococcus aureus* (*S. aureus*) and moderately active against *Escherichia coli* (*E. coli*).

No one of the synthesized compounds had shown activity against *Streptococcus pyogenes* (*S.*

*pyogenes*) and *Pseudomonas aeruginosa* (*P. aeruginosa*).

**Table 1.** The antibacterial activities of synthesized compounds.

Comp. no.	Conc. $\mu\text{g/ml}$	Gram (+)ve			Gram (-)ve		
		<i>S. aureus</i>	<i>S. pyogenes</i>	<i>B. subtilis</i>	<i>E. coli</i>	<i>K. pneumoniae</i>	<i>P. aeruginosa</i>
Zone of inhibition (ZI) in (mm)							
Comp.5	$10^3$	-	-	-	12	-	-
Comp.6	$10^3$	11	-	13	11	11	-
Comp.7	$10^3$	10	-	-	11	10.5	-
Comp.8	$10^3$	9.5	-	-	11	-	-
Amoxicillin	$10^3$	13	5	40	11	11.5	28
Cefixime	$10^3$	16	6	20	22	13	6
Nitrofurantoin	$10^3$	21	9.5	31	16	20	17
Isoniazid	$10^3$	8	-	-	12	-	-
DMSO	Solvent control	-	-	-	-	-	-


(-)= No activity, slightly active (ZI =5-10 mm), moderately active (ZI= 10-15 mm), highly active (ZI= more than 15 mm).<sup>(32,33)</sup>

## Discussion

### Chemistry

**Compound (1)** was esterification product which resulted from reaction between carboxylic acid and an alcohol in the presence of an acid as catalyst. Two stages were involved: **addition** of a nucleophile followed by **elimination** of a leaving

group. Protonation and deprotonation steps also occur during the ester formation which could explain the role of acid in the reaction. Under basic conditions, carboxylate anion will be formed which does not react with an electron-rich nucleophile, so the esterification will be happened in the presence of an acid. Formation of ester is necessary for the success of **step 2** and **5** as explained later.<sup>(34)</sup>



**Figure 2.** Steps of esterification.<sup>(34)</sup>

**Compound (1)** was characterized by carbonyl group of aromatic ester at  $1686\text{cm}^{-1}$  in its FTIR spectrum and  $^1\text{H}$ NMR signals confirmed the presence of  $\text{COOCH}_3$  at 3.80 ppm.

Synthesis of **Compound (2)** and **(5)** is essentially a base catalyzed hydrolysis (hydrazinolysis of ester) which was run under

normal basic condition in which the rate-determining step involves two molecules of hydrazine in which a proton was being transferred




Figure 3. Hydrazinolysis of ester<sup>(35)</sup>.

In case of **compound (2)**; FTIR spectrum was characterized by two stretching vibration bands for the primary amine of hydrazide, at 3310  $\text{cm}^{-1}$  and 3209  $\text{cm}^{-1}$ , respectively, 3256  $\text{cm}^{-1}$  NH amide stretching vibration band, 1628  $\text{cm}^{-1}$  C=O stretching vibration band of amide, and 1601  $\text{cm}^{-1}$   $\text{NH}_2$  bending vibration band.  $^1\text{H}$ NMR signals confirmed the presence of CONH at 9.56 ppm, and  $\text{NH}_2$  at 4.42 ppm.

For **compound (5)**; FTIR spectrum was characterized by asymmetric and symmetric stretching vibration bands of  $\text{NH}_2$  at (3341 and 3217)  $\text{cm}^{-1}$ , respectively, 3256  $\text{cm}^{-1}$  NH amide stretching vibration band, the amide carbonyl stretching vibration band at 1682  $\text{cm}^{-1}$ , and  $\text{NH}_2$  bending stretching vibration band at 1655  $\text{cm}^{-1}$ .  $^1\text{H}$ NMR was revealed the presence of new signals at 9.41 ppm and 4.36 ppm related to CONH and CONHNH<sub>2</sub>, respectively.

Synthesis of **compound (3)** had been carried out by refluxing an ethanolic suspension of **compound (2)** with  $\text{CS}_2$ , in the presence of (KOH) and the cyclization which involved formation of potassium salt of dithiocarbamate as an intermediate could be done through keto form of hydrazide carbonyl or

between them. In the next step, one hydrazine molecule will be left slowly with one molecule of alcohol<sup>(35)</sup>.

enol form. Enol form is the preferred explanation because of the stability of enol form by intramolecular hydrogen bonding.<sup>(36)</sup>




Figure 4. enol form of dithiocarbamate stabilized by intramolecular hydrogen bonding.<sup>(36)</sup>

FTIR spectrum of **compound (3)** was characterized by NH stretching vibration band at 3155  $\text{cm}^{-1}$ , weak SH stretching vibration band at 2669  $\text{cm}^{-1}$ , and in addition to the absence of amide band at 1628  $\text{cm}^{-1}$ .  $^1\text{H}$ NMR was characterized by the appearance of new signal related to SH proton at 14.58 ppm.

**Compound (4)** was obtained by a nucleophilic substitution ( $\text{S}_\text{N}2$ ) reaction between **compound (3)** and ethyl 2-bromoacetate in absolute ethanol and in the presence of anhydrous potassium carbonate as a catalyst.




Figure 5. synthesis of **compound (4)** by nucleophilic substitution ( $\text{S}_\text{N}2$ ) reaction<sup>(37)</sup>.

FTIR spectrum was characterized by 1744  $\text{cm}^{-1}$  stretching vibration band of saturated ester carbonyl.  $^1\text{H}$ NMR was characterized by the presence of new signals at 4.14-4.18 (2H, q, COCH<sub>2</sub>CH<sub>3</sub>), 1.18-1.21 (3H, t, COCH<sub>2</sub>CH<sub>3</sub>), and the absence of SH signal at 14.58 and instead of the appearance of new signal at 4.26 ppm which was related to -SCH<sub>2</sub>.

**Compounds (6-8)** were Schiff base products (imines) which resulted from reaction between aldehydes with a primary amines in mildly acidic conditions and involves six steps; the first three steps produce an intermediate called a carbinolamine and the last three steps convert the carbinolamine into an imine<sup>(38)</sup>.




Figure 6. steps of Schiff base<sup>(38)</sup>.

**FTIR** spectra were characterized by absence of hydrazide  $\text{NH}_2$  asymmetric and symmetric stretching vibration bands; while  $\text{N}=\text{CH}$  imine stretching vibration bands were overlapped with other bands in **FTIR** spectra. **<sup>1</sup>HNMR** were characterized by the appearance of new signals related to  $\text{CONHN}$  which were due to *cis* and *trans* isomers showed 2signals between [11.60,11.63 for **compound (6)**, 11.45,11.48 for **compound (7)**, 11.64,11.69 for **comp.8**], and  $\text{CONHN}=\text{CH}$  which due to *syn/anti-syn* conformers showed 2signals between [7.91, 8.08 for **compound (6)**, 7.86, 8.04 for **compound (7)**, 7.97, 8.14 for **compound (8)**].<sup>(39-42)</sup>

#### Antibacterial activities

Four antibacterial standards were used, amoxicillin to compare *anti-gram (+)ve* activities of the derivatives with it; cefixime to compare *anti-gram(-)ve* activities of the derivatives with it; nitrofurantoin because it is considered hydrazone and contains furan ring which is isoseter with oxadiazole ring, and isoniazid which is hydrazide compound resembles to **compound (5)**. Because **compound (6)** and **compound (7)** are more polar than **compound (8)**, they showed additional activities against *K. pneumoniae* [as polarity increased; there will be extended activity against *gram(-)ve* bacteria, while retained activities against *gram(+)*ve bacteria as in the case of penicillin G and

aminopenicillins].<sup>(43)</sup> Because **Compound (5)** is hydrazide; it is expected to show limited activity against test bacteria and showed agreement with isoniazid.

#### Conclusion:

New oxadiazole derivatives (hydrazide and its Schiff bases), derived from vanillic acid were successfully synthesized by conventional methods. They were characterized and evaluated for their antibacterial activities. **Compound (6)** had shown the broadest spectrum against tested bacteria showed activities against four out of six bacteria. **Compound (7)** had shown moderate activities against *S. aureus*, *E. coli*, and *K. pneumoniae*. **Compound (5)** was moderately and selectively active against *E. coli*. **Compound (8)** was slightly active against *S. aureus* and moderately active against *E. coli*.

#### Acknowledgments

We're grateful to the College of Pharmacy-Department of Pharmaceutical Chemistry-University of Baghdad, for conducting the research, Mr. Ahmed College Ibn -Al-Haitham, for Pure Science, for his polite style and his patience during evaluation of antimicrobial activity, and chemistry analysis center (CAC) staff for their major role in <sup>1</sup>HNMR analysis and evaluation of antibacterial activities of the compounds.

## References

- Al-Mulla A. A review: biological importance of heterocyclic compounds. *Der Pharma Chemica*. 2017;9(13):141-172.
- Balaji K, Bhatt PR, Mallika D, Jha AN. Design, synthesis and antimicrobial evaluation of some mannich base derivative of 2 (2-substituted)-5-amino-thiadiazoles. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2015;7:145-149.
- Khalilullah H, Khan S, Nomani MS, Ahmed B. Synthesis, characterization and antimicrobial activity of benzodioxane ring containing 1, 3, 4-oxadiazole derivatives. *Arabian Journal of Chemistry*. 2016;9:S1029-1035.
- Kaya B, Kaplancıklı ZA, Yurttaş L, Çiftçi GA. Synthesis and biological evaluation of some new pyrimidine bearing 2, 5-disubstituted 1, 3, 4-oxadiazole derivatives as cytotoxic agents. *Turkish Journal of Biochemistry*. 2016;42(2):131-137.
- Dewangan D, T Nakhate K, K Tripathi D, Kashyap P, Dhongde H. Synthesis, Characterization and Screening for Analgesic and Anti-inflammatory Activities of 2, 5-disubstituted 1, 3, 4-oxadiazole Derivatives. *Anti-Inflammatory & Anti-Allergy Agents in Medicinal Chemistry*. 2015;14(2):138-145.
- SUDHA BN, SUBBAIAH NY, SRIKANTH A, REDDY CV, LATHA MS, LAKSHMI PV. Synthesis and Antimicrobial Studies of 5-N-alkyl-1, 3, 4-oxadiazole-2-thiol Derivatives from Fatty Acids. *Asian Journal of Chemistry*. 2017;29(11):2369-2371.
- Yurttaş L, Cantırk Z. The synthesis, antifungal and apoptotic effects of triazole-oxadiazoles against *Candida* species. *European journal of medicinal chemistry*. 2018;144:255-261.
- Ibrahim MK, El-Adl K, Zayed MF, Mahdy HA. Design, synthesis, docking, and biological evaluation of some novel 5-chloro-2-substituted sulfanylbenzoxazole derivatives as anticonvulsant agents. *Medicinal Chemistry Research*. 2015;24(1):99-114.
- Babu RR, Naresh K, Ravi A, Reddy BM, Babu VH. Synthesis of novel isoniazid incorporated styryl quinazolinones as anti-tubercular agents against INH sensitive and MDR M. tuberculosis strains. *Medicinal Chemistry Research*. 2014;23(10):4414-4419.
- Gilani SJ, Khan SA, Alam O, Siddiqui NA. Synthesis and in vitro antimicrobial evaluation of condensed heterocyclic 6-substituted 1, 2, 4-triazolo-[3, 4-b]-1, 3, 4-thiadiazole and 1, 3, 4-oxadiazole derivatives of isoniazid. *Acta Poloniae Pharmaceutica-Drug Research*. 2011;68(2):205-211.
- Popiołek Ł, Biernasiuk A. Hydrazone-hydrazones of 3-methoxybenzoic acid and 4-tert-butylbenzoic acid with promising antibacterial activity against *Bacillus* spp. *Journal of enzyme inhibition and medicinal chemistry*. 2016;31(sup1):62-69.
- Rambabu N, Dubey PK, Ram B, Balram B. Synthesis, characterization and antimicrobial activity of some novel hydrazone derivatives of anacardic acid. *Der Pharma Chemica*. 2015;7(4):90-97.
- Backes GL, Neumann DM, Jursic BS. Synthesis and antifungal activity of substituted salicylaldehyde hydrazones, hydrazides and sulfohydrazides. *Bioorganic & medicinal chemistry*. 2014;22(17):4629-4636.
- Velezheva V., Brennan P., Ivanov P., Kornienko A., Lyubimov S., Kazarian K., et al. Synthesis and antituberculosis activity of indole-pyridine derived hydrazides, hydrazone-hydrazones, and thiosemicarbazones. *Bioorganic & Medicinal Chemistry Letters*. 2015; 26(3):978-985.
- Şenkardeş S, Kaushik-Basu N, Durmaz I, Manvar D, Basu A, Atalay R, et al. Synthesis of novel diflunisal hydrazone-hydrazones as anti-hepatitis C virus agents and hepatocellular carcinoma inhibitors. *European journal of medicinal chemistry*. 2016;108:301-308.
- He H, Wang X, Shi L, Yin W, Yang Z, He H, et al. Synthesis, antitumor activity and mechanism of action of novel 1, 3-thiazole derivatives containing hydrazone-hydrazone and carboxamide moiety. *Bioorganic & Medicinal Chemistry Letters*. 2016;26(14):3263-3270.
- Reddy AK, Kathale NE. Synthesis and Anti-Inflammatory Activity of Hydrazones Bearing Biphenyl Moiety and Vanillin Based Hybrids. *Oriental Journal of Chemistry*. 2017;33(2):971-978.
- Zeeshan S, Naveed M, Khan A, Atiq A, Arif M, Ahmed MN, et al. N-Pyrazoloyl and N-thiopheneacetyl hydrazone of isatin exhibited potent anti-inflammatory and anti-nociceptive properties through suppression of NF-κB, MAPK and oxidative stress signaling in animal models of inflammation. *Inflammation Research*. 2019;68(7):613-632.
- Silverstein RM, Webster FX, Kiemle DJ. Spectrometric identification of organic compounds. 7th ed. Hoboken: John Wiley & Sons. Inc.; 2005. Chapter 2, Infrared Spectrometry; p. 72-126.
- Pavia DL, Lampman GM, Kriz GS, Vyvyan JR. Introduction to spectroscopy 5th ed. United States: Cengage Learning; 2015. Chapter 2, Infrared Spectroscopy; p. 14-106.

21. Pavia DL, Lampman GM, Kriz GS, Vyvyan JR. Introduction to spectroscopy 5th ed. United States: Cengage Learning; 2015. Chapter 5, Nuclear magnetic resonance spectroscopy; p. 215-456.
22. Silverstein RM, Webster FX, Kiemle DJ. Spectrometric identification of organic compounds. 7th ed. Hoboken: John Wiley & Sons. Inc.; 2005. Chapter 3, Proton nmr spectrometry; p. 127-203
23. Hassan OM, Sarsam SW. Synthesis , Characterization and Preliminary Anti-inflammatory Evaluation of New Etodolac Derivatives. 2019;28(1):106–112.
24. Ahmed WS, Razzak Mahmood Kubba AA, Al-Bayati RI. Synthesis and evaluation of antimicrobial activity of new imides and schiff bases derived from Ethyl-4-Amino Benzoate. Oriental Journal of Chemistry. 2018;34(5):2477–2486.
25. Karabanovich G, Zemanová J, Smutný T, Székely R, Šarkan M, Centárová I, et al. Development of 3,5-Dinitrobenzylsulfanyl-1,3,4-oxadiazoles and Thiadiazoles as Selective Antitubercular Agents Active Against Replicating and Nonreplicating Mycobacterium tuberculosis. Journal of Medicinal Chemistry. 2016;59(6):2362–2380.
26. Fadda AA, Abdel-Rahman AH, El-Sayed WA, Zidan TA, Badria FA. Synthesis of novel 1, 3, 4-oxadiazole derivatives and their nucleoside analogs with antioxidant and antitumor activities. Chemistry of Heterocyclic Compounds. 2011;47(7):856-864.
27. Mir I, Siddiqui MT, Comrie AM. Antituberculosis agents. V:  $\alpha$ -[5-(5-nitro-2-furyl)-1, 3, 4-oxadiazol-2-ylthio] acetylhydrazide and related compounds. Journal of pharmaceutical sciences. 1991;80(6):548-550.
28. Han ML, Bekci H, CUMAOĞLU A, KÜÇÜKGÜZEL ŞG. Synthesis and characterization of 1, 2, 4-triazole containing hydrazide-hydrazones derived from (S)-Naproxen as anticancer agents. Marmara Pharmaceutical Journal. 2018;22:559-569.
29. Bauer AW, Kirby WMM, Sherris JC, Truck M. Antibiotic susceptibility testing by standardized single disk method. American Journal of Clinical Pathology. 1966;45(4):493–496.
30. King H. 254. Curare alkaloids. Part IV. Bebeerine and tubocurarine. Orientation of phenolic groups. Journal of the Chemical Society (Resumed). 1939:1157-1164.
31. Kalb L, Groß O. Über die Darstellung von Aldehyden durch Oxydation primärer Säurehydrazide. Berichte der deutschen chemischen Gesellschaft (A and B Series). 1926;59(4):727-737.
32. Dabholkar VV, Gavande RP. Synthesis and antimicrobial activities of novel 1, 4-benzothiazine derivatives. Arabian Journal of Chemistry. 2016;9:S225-229.
33. Ali PS, Meshram JS, Raut RD. Theoretical and synthetic approach towards the biology of some novel monobactam induced sulphonamides: assessing biology through coupling of active ingredients. Jordan Journal of Chemistry. 2011;146(602):1-2.
34. Smith GJ. Organic Chemistry 5th ed. New York: McGraw-Hill, education; 2017. Chapter 22, Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution; p. 868-923.
35. Ali HA. Synthesis, Characterization and Anti-Inflammatory Evaluation of New Potentially Active Naproxen Hydrazones [thesis on the internet]. [Baghdad (Iraq)]: University of Baghdad 2015 [cited 2020 Mar 18]. Available from:[https://www.researchgate.net/publication/287644704\\_Thesis\\_-\\_Synthesis\\_Characterization\\_and\\_Anti-Inflammatory\\_Evaluation\\_of\\_New\\_Potentially\\_Active\\_Naproxen\\_Hydrazones\\_/link/5832c41e08ae138f1c086a4b/download](https://www.researchgate.net/publication/287644704_Thesis_-_Synthesis_Characterization_and_Anti-Inflammatory_Evaluation_of_New_Potentially_Active_Naproxen_Hydrazones_/link/5832c41e08ae138f1c086a4b/download).
36. Shaimaa Abed Saoud. Synthesis of some heterocyclic compounds at position six of 2-methyl phenol and evaluation of their antioxidant properties besides to antibacterial activities for some of them. University of Baghdad ,College of education for pure science/ Ibn-AL-Haitham, Chemistry department, 2017.
37. Karty MJ. Organic Chemistry Principles and Mechanisms. New York-London: W. W. Norton & Company, Inc.; 2014. Chapter 9, Nucleophilic Substitution and Elimination Reactions 1; p. 466-523.
38. Klein D. Organic Chemistry 3rd ed. Hoboken: John Wiley and Sons, Inc.; 2017. Chapter 19, Aldehydes and Ketones; p. 844-897.
39. Kuodis Z, Rutavičius A, Matijoška A, Eicher-Lorka O. Synthesis and isomerism of hydrazones of 2-(5-thioxo-4, 5-dihydro-1, 3, 4-thiadiazol-2-ylthio) acetohydrazide. Open Chemistry. 2007;5(4):996-1006.
40. Dinh NH, Tuyet Lan HT, Thu Trang TT, Van Hoan P. Synthesis and NMR Spectroscopic Characteristics of a Series of Hydrazide-Hydrazones Containing Furoxan Ring Derived from Isoeugenoxycetic Acid. Journal of Heterocyclic Chemistry. 2012;49(4):814-822.
41. Higgs PL, Ruiz-Sanchez AJ, Dalmina M, Horrocks BR, Leach AG, Fulton DA. Enhancing the kinetics of hydrazone exchange processes: an experimental and computational study. Organic & biomolecular chemistry. 2019;17(12):3218-3224.

42. Palla G, Predieri G, Domiano P, Vignali C, Turner W. Conformational behaviour and E/Z isomerization of N-acyl and N-arylohydrazones. *Tetrahedron*. 1986;42(13):3649-3654.
43. Beale JM, Block JH. Wilson and Gisvold's Textbook of organic medicinal and pharmaceutical chemistry. 12th ed. Philadelphia: Lippincott Williams & Wilkins, a Wolters Kluwer business; 2011. Chapter 8, Antibacterial Antibiotics; p. 258-329.

