

Association of Oxidative Stress Markers with Cholelithiasis

Omer S.Sadiem ^{*1}, Mohamed A. Taher ^{**} and Seenaa S. Amin ^{**}

* Department of Clinical Laboratory Sciences, College of Pharmacy, Tikrit University, Tikrit, Iraq.

** Department of Clinical Laboratory Sciences, College of Pharmacy, University of Baghdad, Baghdad, Iraq.

Abstract

Oxidative stress markers are of important diagnostic parameters for many disorders including cholelithiasis. This present study has aimed to assess the state of oxidative stress in symptomatic radiographically confirmed (Cholelithiasis) patients by measuring two parameters used as oxidative stress parameters which are serum myeloperoxidase (MPO) and superoxide dismutase (SOD). This study was carried out on 100 patient diagnosed as (Cholelithiasis) patients with 30 age and sex matched healthy controls by measuring serum (MPO) and (SOD) by ELIZA technique. Results showed significantly decrease in antioxidant enzyme(SOD) and increase in serum level of (MPO) comparing with controls.

Keywords: Cholelithiasis , Oxidative stress, MPO, SOD.

مصاحبة عوامل الاجهاد التاكسدي مع حصوة المرارة
عمر صلاح صديم^{1*}، محمد عباس طاهر^{**} و سناء صادق امين^{**}

* فرع العلوم المختبرية السريرية، كلية الصيدلة، جامعة تكريت، تكريت، العراق.
** فرع العلوم المختبرية السريرية، كلية الصيدلة، جامعة بغداد، بغداد، العراق.

الخلاصة

عوامل الاجهاد التاكسدي هي مؤشرات قوية للعديد من الحالات المرضية بما في ذلك (حصوة المرارة). البيانات الموجودة على العلاقة بين حصوة المرارة وعلامات الاجهاد التاكسدي مخفزة ولكن محدودة. لذلك هدفت الدراسة الحالية الى تقييم حالة الاجهاد التاكسدي لدى مرضى حصوة المرارة المشخصين بالاشعة السينية من خلال عاملين من عوامل الاجهاد التاكسدي (MPO & SOD). شملت الدراسة 100 مريض مشخصين بحصوة المرارة مع 30 من الاصحاء كمجموعة سيطرة مطابقة لمجموعة المرضى في العمر والجنس، وظهرت النتائج انخفاض معنوي بمستوى انزيم SOD مع ارتفاع معنوي بمستوى MPO بالمقارنة مع مجموعة السيطرة.

الكلمات المفتاحية: حصوة المرارة، الاجهاد التاكسدي، MPO، SOD.

Introduction

Cholelithiasis is a worldwide disease and it remains to be one of the most common health problems leading to surgical intervention ^(1,2). Cholelithiasis is common with the incidence ranging from (10-20)% of the world population ⁽³⁾, the prevalence of cholelithiasis varies considerably within population; the highest known prevalence is among American Indians with up to (60-70) % in females ⁽⁴⁾ and (10-15) % in white adults of developed countries ⁽⁵⁾. Cholelithiasis is classified as either cholesterol stones (Cs), pigment stones (Ps), or mixed stones (Ms)⁽⁶⁾. These categories of gallstones can be identified according to their predominant chemical composition ⁽⁷⁾. Two thirds of individuals with cholelithiasis is asymptomatic ⁽⁸⁾. In these patients, the reasons why symptoms develop are unknown ⁽⁹⁾. Of these, symptoms will develop in 1% to 4% per year ^(10, 11). Oxidative stress (OS) is a disturbance in the oxidant-antioxidant balance leading to potential cellular damage. Most cells can tolerate a mild degree of oxidative stress, because they have sufficient antioxidant

defense capacity and repair systems, which recognize and remove molecules damaged by oxidation. The imbalance can result from a lack of antioxidant capacity caused by disturbances in production and distribution, or by an overabundance of reactive oxygen species (ROS) from other factors ⁽¹²⁾. Increased levels of an accelerated generation of reactive oxygen species and toxic degradation products of lipid peroxidation have been reported in the serum of individuals with cholelithiasis⁽¹³⁾. It is well known that, in chronic diseases such as cholelithiasis, the active inflammatory response is induced with neutrophilic infiltration. These neutrophils, macrophages and/or monocytes produce ROS which may cause DNA damage to the adjacent cells ^(14, 15). Superoxide dismutase or SOD is an enzyme that in humans is encoded by the SOD1 gene, located on chromosome 21. SOD is one of three human superoxide dismutase ^(16, 17). Superoxide dismutase (SOD) is an enzyme that in humans is encoded by the SOD1 gene, located on chromosome 21. SOD is one of three human superoxide dismutase ^(16, 17).

¹Corresponding author E-mail: Omer S. Sadiem: omer_salah82@gmail.com

Received: 26 /1/2014

Accepted: 12/10/2014

Myeloperoxidase (MPO) is a peroxidase enzyme that in humans is encoded by the MPO gene⁽¹⁸⁾. Myeloperoxidase is most abundantly expressed in neutrophil granulocytes (a subtype of white blood cells). It is a lysosomal protein stored in azurophilic granules of the neutrophil. MPO has a heme pigment, which causes its green color in secretions rich in neutrophils, such as pus and some forms of mucus⁽¹⁹⁾. Gallstones and kidney stones are diseases with high prevalence⁽²⁰⁾. There has been a marked increase in the prevalence and incidence of these stones within the last 22 years⁽²¹⁾. In spite of different composition and different locations of their formation both types of stones may be associated⁽²²⁾. Obesity, insulin resistance and women with postmenopausal hormone use are at risk factors for renal stones as well as gallstones^(23, 24).

Materials and Methods

Subjects

This study was conducted at Tikrit Teaching Hospital, Tikrit, Iraq, and the laboratory investigations were done at Tikrit University, Department of Central Laboratories. It is carried from December 2012 to June 2013. A total of 100 patients (group A) were included in the study, among which 68 were females patients and 32 were males in the age group of (20-80) years and mean age (48.38 ± 1.17), to compare the results of 30 (age and sex matched) healthy controls (Group B) (18 females, 12 males) with mean age (47.16 ± 1.5) years were also included.

Patient's including criteria:

All patients were scanned for presence of gallstone by an abdominal ultrasonography (the standard diagnostic test for gallstone), and to confirm the absence of kidney stone.

Patient's exclusion criteria:

1. Detection complicated gallstone diseases like (acute cholecystitis, pancreatitis).
2. Patients with liver cirrhosis, viral hepatitis, renal failure and thyroid diseases and any type of cancer.
3. Patients with a history of alcohol consumption and smoking.
4. Subjects suffered from diseases like asthma.

Questionnaire

Subjects were asked to complete a questionnaire that asked for information on name, sex, age, marital state, number of children, pregnancy (for females only), family history of GS, chronic diseases, urinary tract complications, weight, height, occupation, residence, smoking habits, alcohol consumption, and drugs used.

Ultrasound for patients with cholelithiasis and nephrolithiasis were checked to confirm the

presence of calculi, number of stones and their average diameter.

All the laboratory investigations were done in Tikrit University, Central Laboratories Department.

Sample collection

After 12 hours of fasting, venous blood samples, about 10 ml were collected from patients before laparoscopic cholecystectomy and from healthy volunteers in plain tubes. After allowing the blood to clot at room temperature for about 15-30 min, blood samples were centrifuged at 3000 rpm for about 15-30 min to obtain serum. The serum were separated in multiple epiendroff tubes (0.5 ml) to prevent thawing of the serum frequently, The serum was stored and frozen at -20°C until analysis was performed.

Clinical and laboratory evaluation

1. Determination of Myeloperoxidase (MPO):

The test is based on the immobilisation of highly purified MPO to the solid phase of microtiter strips and subsequent binding of anti-MPO antibodies from patient serum by using ELISA technique. The bound antibodies are detected with a peroxidase-labeled secondary antibody that is directed against human IgG. After addition of substrate solution, a color appears which intensity is proportional to the concentration and/or the avidity of the detected antibodies. Following the addition of stop solution, the colour switches from blue to yellow. The concentration of MPO in the samples is then determined by comparing the optical density of the samples to the standard curve.⁽²⁵⁾

2. Determination of human superoxide dismutase [Cu-Zn] (SOD1):

This assay employs the quantitative sandwich enzyme immunoassay technique (ELISA). Antibody specific for SOD1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any SOD1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for SOD1 is added to the wells. After washing, avidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of SOD1 bound in the initial step. The color development is stopped and the intensity of the color is measured⁽²⁶⁾.

Statistical analysis

The results were expressed as mean \pm standard error of mean (SEM). Student's t-test was used to examine the degree of significance. P values less than 0.05 was considered significant. The statistical analysis was performed using the Statistical Package for Social Sciences (SPSS 17).

Table (1): Demographic presentation of Cholelithiasis patients and control.

Parameter	Control N =30	Patient N = 100	P- value
Age (mean \pm SD)	48.38 \pm 1.17	47.16 \pm 1.5	0.724
F : M	18:12 (60%)	32:68 (68%)	0.300
Pregnancy	2 (6.6%)	4 (5.8%)	0.485
Family History	10 (34%)	55 (55%)	0.092
H.T	4 (4%)	14 (14%)	0.748
D.M	3 (10%)	5 (5%)	0.748

Continuous variables presented as Mean \pm SD, and discrete variables as numbers and frequencies; F:M= female and male ratio; n= Number ; H.T= hyper tension; D.M= Diabetes mellitus; (*) = significant difference P < 0.05

Results

Demographic presentation of 100 cholelithiasis patients and 30 controls were elucidated in (table 1). Frequency matching was successful with Patient and control groups having similar age distributions (mean \pm SD) ages.

Table 2 showed the result of the parameters for both controls and cholelithiasis patients.

The current study demonstrates a significant difference in SOD serum level between controls and patients P-value (0.026). so, there was a significant decrease in the serum levels of SOD indicates decreasing in antioxidant activity. While for MPO, current study shows also high significant difference between patient and control P-value (0.001).

Table (2):Parameters of the patients and controls

Parameter s	Control N = 30	Patients N = 100	P- value
SOD	0.67 \pm 0.51	0.42 \pm 0.29	0.026*
MPO	7.01 \pm 1.55	9.27 \pm 2.66	0.001*

Value were presented as mean \pm SD of mean; (*) = significance difference, (P<0.05)
N= no. of individuals in each group.

Discussion

Similar finding were shown by Correa P.⁽²⁷⁾ and ByungKyu Park *et al.*⁽²⁸⁾, they indicate significantly decrease in SOD serum levels in patient diagnosed with many gastrointestinal diseases like acute pancreatitis gallbladder cancer and Cholelithiasis. Also the current study supported by the results of Kaur T. and Kaur S.2010⁽²⁹⁾ found "significant decrease in antioxidant enzymes activities (superoxide dismutase) in patient with gallstone comparing with controls". Also RuhiDixit, *et al.*⁽³⁰⁾ and AlpaslanTerzet *al.*⁽³¹⁾ have the same findings. Gallstones can induce inflammation in the gallbladder wall, and the composition of bile changes, at the same time the bilirubin metabolism, which is a potent antioxidant by radical scavenging and reducing activities, may be altered⁽³²⁾. The changes in bile composition can increase the biliary free radical formation⁽³³⁾. Increased levels of inducible nitric oxide (NO) synthesis activity was shown in inflamed gallbladders which has an effect on elevation of oxidative stress and on fluid transport as well⁽³⁴⁾. Moreover, inflammatory changes of the gallbladder mucosa are associated with granulocyte infiltration; the activated phagocytes can produce reactive oxygen metabolites, and lead to oxidative stress. Free radicals and other peroxides derivatives produced physiologically in the human body and increased in many pathological conditions, diffuse into the blood⁽³⁵⁾. Here, antioxidant components of plasma overwhelm them and they are consumed. Total antioxidants; therefore, is detected as being significantly lower in the cholelithiasis patients than in the controls. Many antioxidants and oxidants were studied to determine their effects. In particular, vitamin C, which is a strong external antioxidant, serum ascorbic acid level was inversely related to prevalence of gallbladder disease⁽³⁶⁾.

Bilirubin free radicals can damage the liver cells, which can induce the change of ingredients of the bile by decreasing the amount of bile acids. The abnormal metabolism in hepatocyte can lead to hydrolysis of the conjugated bilirubin, increase in the concentration of free bilirubin, which in turns make bilirubin supersaturated to the bile, and promote the formation of pigment gallstone, during the gallstone formation, bilirubin reacts with the active-oxygen species formed in vivo followed by its polymerization, aggregation, and calcification. It is speculated to be an important step in the formation of pigment gallstones⁽³⁷⁾.

For the MPO, similar findings of this study were found by (H.Ojima *et al.* 2005) who

consider serum level of MPO as diagnostic parameter for undifferentiated carcinoma and small cell carcinoma in gallbladder such as Extramedullary myeloid tumour (EMMT) ⁽³⁸⁾, also (Alfredo Menendez 2009) with (ShikoKuribaYashi , *et.al.* 2004) agree with this finding and used MPO and other oxidative stress markers as effective diagnostic parameters for many bladder dysfunction including cholelithiasis^(39,40).

Also in animal models, Rege RV, Prystowsky JB ⁽⁴¹⁾ found a significant elevation in both MPO and IL-1 in cholelithiatic animals, and (ÖzgürKasımayet *al*)2006 who found significant increase in MPO level in gastrointestinal diseases including Cholelithiasis⁽⁴²⁾.

Conclusion

Cholelithiasis patients showed low antioxidant activity, and high MPO serum levels .as shown in this study; the oxidative stress markers are useful in the pathogenesis and diagnosis of Cholelithiasis.

Recommendation

1. Measuring more parameters that used in the diagnoses of cholelithiasis.
2. Measuring more oxidative stress markers and antioxidant enzymes in cholelithiasis patients.

References

1. Charles F. Brunicaardi: Shwartz principle of surgery. 9th edition , McGraw - Hill companies, U.S. 2010 pp: 1136-1161.
2. Sandler RS, Everhart JE, Donowitz M, et al: The burden of selected digestive diseases in the United States. *Gastroenterology* 2002; 122:1500–11.
3. Pradhan SB1, Joshi MR2, Vaidya A: Prevalence of different types of gallstone in the patients with cholelithiasis at Kathmandu Medical College, Nepal. *Kathmandu University Medical Journal* 2009; 7(3): 268-271.
4. James S. Dolly Anna SF: *Sherlocks Disease of the Liver And Biliary System*. 12th ed. 2011. Blackwell Publishers P264.
5. Stinton Laura, Meyers M, et al: Epidemiology of Gallstone. *Clinics of North America* 2010; 39(2): 157-169.
6. Novacek G. Gender and gallstone disease. *Wien Med Wochenschr.* 2006; 156(19):527-533.
7. Vanberge-Henegouwen GP, Venneman NG, Van Erpecum KJ, Portincasa P: Drugs affecting biliary lipid secretion and gallbladder motility: their potential role in gallstone treatment and prevention. *Curr Drug Targets Immune Endocr Metabol Disord.* 2005; 5(2): 185-191.
8. Sidney Cohen MD, Bruce R. , Bacon MD, et al : National Institutes of Health State-of-the-Science Conference Statement: ERCP for diagnosis and therapy, January 14-16, 2002. *Gastrointestinal Endoscopy* 2002; 56(6):803-809.
9. Cynthia W. Ko, MS, Sum P. Lee: Epidemiology and natural history of common bile duct stones and prediction of disease. *Gastrointestinal Endoscopy.* 2002; 56(6): S165-S169
10. Halldestam I, Enell EL, Kullman E, Borch K: Development of symptoms and complications in individuals with asymptomatic gallstones. *Br J Surg* 2004; 91: 734–738.
11. National Institutes of Health. NIH Consensus Statement on Gallstones and Laparoscopic Cholecystectomy 1992; 10 (3):1-20.
12. Hye-Lin Ha, Hye-Jun Shin, Mark A Feitelson, et al: Oxidative stress and antioxidants in hepatic pathogenesis. *World J Gastroenterol* 2010 ; 16(48): 6035-6043.
13. Geetha A. Evidence for oxidative stress in the gall bladder mucosa of gallstone patients. *J BiochemMolBiolBiophys* 2002; 6:427-32.
14. Matsui A, Ikeda T, Enomoto K, et al: Increased formation of oxidative DNA damage, 8-hydroxy-2'-deoxyguanosine, in human breast cancer tissue and its relationship to GSTP1 and COMT genotypes. *Cancer Lett.* 2000; 151:87-95.
15. Galli F, Piroddi M, Annetti C, et al: Oxidative stress and reactive oxygen species. *Contrib Nephrol* 2005; 149:240-60.
16. Milani P, Gagliardi S, Cova E, Cereda C (2011). "SOD1 Transcriptional and Posttranscriptional Regulation and Its Potential Implications in ALS.". *Neurol Res Int.* 2011: 458427. Doi : 10.1155/2011/458427. PMC 3096450. PMID 21603028
17. Rosen DR, Siddique T, Patterson D, Figlewicz DA, Sapp P, Hentati A, Donaldson D, Goto J, O'Regan JP, Deng HX. "Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis". *Nature* 1993; 362 (6415): 59–62.
18. Blair-Johnson M, Fiedler T, Fenna R. "Human myeloperoxidase: structure of a cyanide complex and its interaction with bromide and thiocyanate substrates at 1.9 Å resolutions". *Biochemistry* 2001; 40 (46): 13990–7.

19. Klebanoff SJ . "Myeloperoxidase: friend and foe". *J. Leukoc. Biol.* 2005; 77 (5): 598–625.
20. Cornelia Weikert , Steffen Weikert, Matthias B. Schulze, Tobias Pischon, et al: Presence of Gallstones or Kidney Stones and Risk of Type 2 Diabetes. *Am. J. Epidemiol.* 2010 ;171 (4): 447-454.
21. A. Hesse, E. Brandeb, D. Wilbert, K.-U. Köhrmann, P. Alkand: Study on the Prevalence and Incidence of Urolithiasis in Germany Comparing the Years 1979 vs. 2000. *European Urology* 2003;44(6):709–713.
22. Eric N. Taylor, Andrew T. Chan: Cholelithiasis and the Risk of Nephrolithiasis. *Journal of Urology.* 2011;186(5): 1882-1887.
23. Domenico R, Giuseppe M, et al: Association between metabolic syndrome and nephrolithiasis in an inpatient population in southern Italy: role of gender, hypertension and abdominal obesity. *Nephrol Dial Transplant.* 2009;24(3):900-906.
24. Maalouf NM, Alicia H. Sato, Brian J. Welch : Postmenopausal Hormone Use and the Risk of Nephrolithiasis: Results From the Women's Health Initiative Hormone Therapy Trials. *Arch Intern Med.* 2010;170(18):1678-1685.
25. Milla,C.; Yang,S.; Cornfield,D.N.; Brennan,M.L.; Hazen,S.L.; Panoskaltis-Mortari,A.; Blazar,B.R.; Haddad,I.Y.. Myeloperoxidase deficiency enhances inflammation after allogeneic marrow transplantation. *Am.J.Physiol Lung Cell Mol.Physiol* 2004;287:L706-L714.
26. Mount,J.N.*Clin.pathology* 1986; 22,12
27. Correa P. The role of antioxidants in gastric carcinogenesis. *Crit Rev Food Sci Nutr* 1995; 35: 59-64
28. ByungKyu Park, Jae Bock Chung et.al Role of oxygen free radicals in patients with acute pancreatitis *World J Gastroenterol* 2003;9(10):2266-2269
29. Kaur T, Kaur S.: Pathophysiological conditions in cholelithiasis formation in North Indian population: spectroscopic, biophysical, and biochemical study. *Biol Trace Elem Res.* 2010; 138(1-3):79-89.
30. Dixit et al.: Chromosomal structural analysis in carcinoma of the gallbladder. *World Journal of Surgical Oncology* 2012; 10:198
31. AlpaslanTerz et al: Oxidative Status and Plasma Prolidase Activity in Patients with Gallstones. *TrakyaUniv Tip FakDerg* 2010;27(4):358-362
32. Sipos P, Krisztina H, Blázovics A, et al: Cholecystitis, gallstones and free radical reactions in human gallbladder. *MedSciMonit.* 2001;7:84-8.
33. Chamulitrat W, Carnal J, Reed NM: In vivo endotoxin enhances biliary ethanol-dependent free radical generation. *Am J Physiol*, 1998;274: 653-61.
34. Nilsson B, Delbro D, Hedin L, et al: Role of nitric oxide induction of inflammatory fluid secretion by the mucosa of the feline gallbladder. *Gastroenterology*, 1996;110: 598-606.
35. Sipos P, Krisztina H, Blázovics A, et al: Cholecystitis, gallstones and free radical reactions in human gallbladder. *MedSciMonit.* 2001;7:84-8.
36. Joel A. Simon, Esther S. Hudes: Serum Ascorbic Acid And Gallbladder Disease Prevalence Among US Adults. *ARCH INTERN MED* 2000; 160: 931-936.
37. Liu X-T, Hu J : Relationship between bilirubin free radical and formation of pigment gallstone. *World J Gastroenterol* 2002; 8(3):413–417.
38. H Ojima, T Hasegawa, Y Matsuno, and M Sakamoto: Extramedullary myeloid tumour (EMMT) of the gallbladder, *J ClinPathol.* 2005; 58(2): 211–213
39. Alfredo Menendez, Ellen T. Arena, et,al : Salmonella Infection of Gallbladder Epithelial Cells Drives Local Inflammation and Injury in a Model of Acute Typhoid Fever in the journal of infectious diseases 2009; 200 (11).
40. ShikoKuribYashi, Tsuyoshi Monden , et al: Usefulness of Cholescintigraphy with Lipid Meal Loading for Diagnosis and Determination of Cholecystectomy in a Patient with Gallbladder Dysfunction. *The journal of Japanese society of internal medicine*2004; 43 (5): 393-396 .
41. Rege RV, Prystowsky JB: Inflammatory properties of bile from dogs with pigment gallstones. *Am J Surg:* 1996;171(1):197-201.
42. ÖzgürKasımay, EsraGüzel, Ali Gemici, Asead Abdyli.et al: Colitis-induced oxidative damage of the colon and skeletal muscle is ameliorated by regular exercise in rats: the anxiolytic role of exercise. *Physiology in Press;* 2006.